



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2021

On Multifaceted Definitions of Multilevel Societies: Response to Papageorgiou and Farine

Grueter, Cyril C ; Qi, Xiaoguang ; Zinner, Dietmar ; Bergman, Thore ; Li, Ming ; Xiang, Zuofu ; Zhu, Pingfen ; Miller, Alex ; Krützen, Michael ; Fischer, Julia ; Rubenstein, Daniel I ; Vidya, T N C ; Li, Baoguo ; Cantor, Maurício ; Swedell, Larissa

DOI: <https://doi.org/10.1016/j.tree.2020.10.012>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-197199>

Journal Article

Accepted Version

Originally published at:

Grueter, Cyril C; Qi, Xiaoguang; Zinner, Dietmar; Bergman, Thore; Li, Ming; Xiang, Zuofu; Zhu, Pingfen; Miller, Alex; Krützen, Michael; Fischer, Julia; Rubenstein, Daniel I; Vidya, T N C; Li, Baoguo; Cantor, Maurício; Swedell, Larissa (2021). On Multifaceted Definitions of Multilevel Societies: Response to Papageorgiou and Farine. *Trends in Ecology & Evolution*, 36(1):17-19.

DOI: <https://doi.org/10.1016/j.tree.2020.10.012>

On multifaceted definitions of multilevel societies: Response to *Papageorgiou and Farine*

Cyril C. Grueter*, Xiaoguang Qi, Dietmar Zinner, Thore Bergman, Ming Li, Zuofu Xiang,
Pingfen Zhu, Alex Miller, Michael Krützen, Julia Fischer, Daniel I. Rubenstein, T. N. C. Vidya,
Baoguo Li, Maurício Cantor and Larissa Swedell

CCG: School of Human Sciences, The University of Western Australia, Perth, WA 6009,
Australia

CCG: Centre for Evolutionary Biology, School of Biological Sciences, The University of Western
Australia, Perth, WA 6009, Australia

CCG: International Centre of Biodiversity and Primate Conservation, Dali University, Dali,
Yunnan 671003, China

XQ: Shaanxi Key Laboratory for Animal Conservation, College of Life Sciences, Northwest
University, Xi'an, 710069, China

DZ: Cognitive Ethology Laboratory, German Primate Center (DPZ), Leibniz Institute for
Primate Research, 37077 Göttingen, Germany

DZ: Leibniz ScienceCampus for Primate Cognition, 37077 Göttingen, Germany

TB: Department of Psychology, University of Michigan, Ann Arbor, USA

TB: Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, USA

ML: CAS Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology,
Chaoyang District, Beijing 100101, China

ML: Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences,
Kunming 650223, China

ZX: College of Life Science and Technology, Central South University of Forestry and
Technology, Changsha, Hunan 410004, China

PZ: CAS Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology,
Chaoyang District, Beijing 100101, China

AM: School of Human Sciences, The University of Western Australia, Perth, WA 6009, Australia

MK: Department of Anthropology, University of Zurich, 8057, Zürich, Switzerland

JF: Cognitive Ethology Laboratory, German Primate Center (DPZ), Leibniz Institute for
Primate Research, Kellnerweg 4, 37077 Göttingen, Germany

JF: Department for Primate Cognition, Georg-August-University of Göttingen, Göttingen,
Germany

DR: Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA

TNCV: Evolutionary and Organismal Biology Unit, Jawaharlal Nehru Centre for Advanced Scientific Research (JNCASR), Jakkur, Bengaluru 560064, India

BL: Shaanxi Key Laboratory for Animal Conservation, College of Life Sciences, Northwest University, Xi'an, 710069, China

BL: Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Science, Kunming, 650223, China

MC: Department for the Ecology of Animal Societies, Max Planck Institute of Animal Behavior, Konstanz, 78464, Germany

MC: Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, 78464, Germany

MC: Department of Biology, University of Konstanz, Konstanz, 78464 Germany

MC: Departamento de Ecologia e Zoologia, Universidade Federal de Santa Catarina, Florianópolis, 88048-970, Brazil

MC: Centro de Estudos do Mar, Universidade Federal do Paraná, Pontal do Paraná, 83255-000, Brazil

MC: School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Johannesburg, 2000, South Africa

LS: Department of Anthropology, Queens College, City University of New York, Flushing, NY 11367-1597, USA

LS: New York Consortium in Evolutionary Primatology, New York, NY, USA

LS: Anthropology, Biology and Psychology Programs, CUNY Graduate Center, 365 Fifth Avenue, New York, NY 10016, USA

LS: Department of Archaeology, University of Cape Town, Rondebosch 7701, Cape Town, South Africa

Papageorgiou and Farine [1], in their comment on our recent synthesis of animal multilevel societies [2], provide several examples of nestedness in avian social systems and call for a fuller incorporation of birds into our theoretical framework. We focused mainly on mammals to construct our proposed framework because multilevel societies are best known from this taxonomic group. Papageorgiou and Farine [1] point out several bird species that form nested social arrangements and argue that, by diving deeply into examples from birds, there may be variations in form that meet our criteria for a multilevel society—a social system with a stable core level and at least one recognizable upper level. Papageorgiou and Farine [1] raise two questions: first, whether the operational definition of multilevel societies should be relaxed to accommodate bird species that show more stability at higher levels of society; and second, whether societies with multiple tiers resulting solely from habitat preferences instead of social preferences should be classified as multilevel societies.

With regard to the first question, Papageorgiou and Farine [1] argue that our criteria should be relaxed to include social stability at any level of a nested society and that the core unit does not necessarily need to be the lowest level. However, we maintain that (i) encompassing every society with a demonstrable modular structure would invalidate the inclusion criteria we developed for distinguishing multilevel societies from other types of systems with detectable substructure; and (ii) a more permissive definition would hamper efforts in identifying the eco-evolutionary drivers of multilevel societies *sensu stricto*. Moreover, maintaining this distinction between multilevel societies with at least two consistent levels and societies with a single stable level combined with other unstable associations (as seen in several bird taxa) is crucial because evolutionary processes such as information flow and disease transmission are expected to differ between these types of societies. That said, we are not opposed to the idea that the most stable levels can be higher levels rather than the core unit. Societies with stable intermediate and upper levels, yet unstable lower levels (as exemplified by vulturine guineafowl *Acryllium vulturinum*; [3]), could constitute multilevel societies, as we originally acknowledged [2]. In fact, in our discussion of the multilevel alliance system of the Shark Bay bottlenose dolphins (*Tursiops* sp.) we pointed out that this system deviates from our definition in that the highly cohesive and stable units (second-order alliances) occur at a level above the core unit. According to our framework, however, a society with unstable core units, no intermediate level, and clearest membership at a higher level would not represent a multilevel society but instead a system with atomistic fission-fusion dynamics. This is exemplified by the northern muriqui (*Brachyteles hypoxanthus*), which exhibits flexible association patterns at basal levels but cohesiveness at the upper level, both spatially and temporally [4].

Papageorgiou and Farine [1] shortlist five possible avian candidates for multilevel societies. But which ones do actually fit our definition? Beyond vulturine guineafowls, bell miners (*Manorina melanophrys*) may be a good fit, as discrete social organization manifests itself on at least three levels [5]. Two additional species appear to ‘tick the boxes’. In spectacled parrotlets (*Forpus conspicillatus*), monogamous pairs are embedded within putatively stable groups which then form flocks [6]. This does appear to be a multilevel society, but more data on association patterns are probably needed to rule out alternatives. In white-fronted bee-eaters (*Merops bullockoides*), there is an intermediate social tier between the breeding pair and the larger colony [7], which could qualify their society as multilevel. Careful screening of the literature will likely reveal similar systems in hitherto neglected taxa.

The second major question raised by Papageorgiou and Farine [1] pertains to whether the definition of multilevel societies should include structured societies brought about by shared spatial preferences or other ecological opportunities (e.g., preferences for certain sleeping and foraging sites) instead of social preferences. In some cases it is not known to what degree the formation of a distinct level is ecologically induced or the result of individual expressions of social preference. For some species, we still do not know exactly where the social glue ends and habitat-induced overlap begins, i.e. at which spatial scales the social processes are paramount. While core (and intermediate) levels are undoubtedly genuine social groupings that crystallize for reproductive purposes and social support, we should be open to the idea that assortment into upper-level groupings is not exclusively driven by social preferences but may also involve ecologically driven mechanisms. Apex levels in particular often represent aggregations of individuals resulting entirely from an external factor, e.g. the magnetic effects that localized resources exert on social units, as is the case with rare safe sleeping sites in the classic example of the troop level in hamadryas baboons [8]) which are very different from a ‘group’ of individuals in which the presence and identity of conspecifics matter. Lack of individualization, however, does not preclude the possibility that individuals derive benefits from being associated with the apex grouping level (e.g. ‘safety in numbers’).

Because we cannot always ascertain whether a particular level of a multilevel society is socially or ecologically driven, this should not be a decisive factor in classifying a society as multilevel or not. In our view, the critical criteria are (1) consistency of individual membership in each level over time and (2) spatio-temporal cohesion of the core and upper levels. An important goal for future research is to quantify the relative contributions of social processes and ecological factors in shaping additional levels in animal societies. This will require fine-grained data on how animals

move relative to each other, and, ideally, on how they may perceive each other as individuals and members of distinct social levels. It would also be interesting to compare the consistency of groupings that are purportedly socially versus ecologically driven. Papageorgiou and Farine [1] argue that studies from birds can allow social versus non-social drivers of nested social levels to be disentangled. While we agree that a broader perspective is useful, it is not clear to us how birds are in any way more suitable than mammals for distinguishing the drivers of social levels, particularly as few of their examples are unambiguously multilevel. Clearly, much further work is needed, and data from mammals, birds and other taxa will be crucial in enriching and refining our understanding of the evolutionary processes responsible for the emergence and maintenance of this intriguing social system.

1. Papageorgiou, D. and Farine, D. (in press) Multilevel societies in birds. *Trends in Ecology & Evolution*.
2. Grueter, C.C. et al. (2020) Multilevel organisation of animal sociality. *Trends in Ecology & Evolution* 35 (9).
3. Papageorgiou, D. et al. (2019) The multilevel society of a small-brained bird. *Current Biology* 29, R1120-R1121.
4. Strier, K.B. et al. (1993) Demography and social structure of one group of marmosets (*Brachyteles arachnoides*). *International Journal of Primatology* 14, 513-526.
5. Painter, J.N. et al. (2000) Complex social organization reflects genetic structure and relatedness in the cooperatively breeding bell miner, *Manorina melanophrys*. *Molecular Ecology* 9, 1339-1347.
6. Wanker, R. et al. (1998) Discrimination of different social companions in spectacled parrotlets (*Forpus conspicillatus*): evidence for individual vocal recognition. *Behavioral Ecology and Sociobiology* 43, 197–202.
7. Hegner, R.E. et al. (1982) Spatial organization of the white-fronted bee-eater. *Nature* 298, 264-266.
8. Kummer, H. (1968) *Social Organization of Hamadryas Baboons: A Field Study*, The University of Chicago Press.